

Physical/Occupational Therapy Prescription

Name: Date of Birth: Diagnosis: Ulnar Collateral Ligament Insufficiency Code: S53.39 Procedure: UCL Repair with Internal Brace (Tommy John Surgery)
Surgery Date:
Instructions:
Weeks 2-4 Splint immobilization until two weeks post-operatively. Work to regain full range of motion by 4 weeks post-operatively. Protect the still healing ulnar ligament by avoiding elbow valgus torque activities. Emphasize maintenance of flexibility and strength of fingers, shoulder, scapula, core, and legs.
Weeks 4-6 - Being isotonic strengthening at the elbow. Stretching and strengthening of the periscapular stabilizers, rotator cuff, deltoid, legs, and core. Initiate the Thrower's Ten Exercise program.
Weeks 6-12: Continue to progress stretching and flexibility. Progress to Advanced Thrower's Ten, advance weights/bands. Begin isotonic lifting program including bench press, seated row, lat pull-downs, triceps push downs, and biceps curls Begin plyometrics (specifically bouncing a medicine ball into a trampoline), progressing from 2- handed (6-8 weeks) to 1-handed (8-10 weeks).
>12 weeks: Initiate progressive interval throwing program. Initiate progressive interval hitting program. Continue strengthening and stretching and Advanced Thrower's Ten program.
Please emphasize a home exercise program.
Criteria for return to competitive throwing: completion of interval throwing program, greater strength in the shoulder on the operative arm than the non-operative arm.
Modalities: Heat before and ice after therapy. Massage prn
Frequency: 2-3 times/week
Duration: 6 weeks
Signature:
Date: